



Soils Map



Soils data provided by USDA and NRCS.



State: **Michigan**
 County: **Eaton**
 Location: **1-1N-3W**
 Township: **Hamlin**
 Acres: **256.72**
 Date: **1/3/2018**



Area Symbol: MI045, Soil Area Version: 14

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
MaB	Marlette loam, 2 to 6 percent slopes	87.23	34.0%		4.5ft.	Moderately well drained	Ile						
MaC	Filer loam, 6 to 12 percent slopes	34.59	13.5%		> 6.5ft.	Well drained	IIle						
BoB	Boyer sandy loam, 0 to 6 percent slopes	34.43	13.4%		> 6.5ft.	Well drained	IIIs	3.8	80	14	60	30	35
BnB	Boyer loamy sand, 0 to 6 percent slopes	21.05	8.2%		> 6.5ft.	Well drained	IIIs	3.8	80	14	60	30	35
WaA	Wasepi sandy loam, 0 to 3 percent slopes	20.60	8.0%		1.5ft.	Somewhat poorly drained	IIIs	3.8	85	14	65	32	35
OwB	Owosso-Marlette sandy loams, 1 to 6 percent slopes	14.42	5.6%		> 6.5ft.	Well drained	Ile	5	115	19	95	35	55
BoC	Boyer sandy loam, 6 to 12 percent slopes	9.58	3.7%		> 6.5ft.	Well drained	IIle	3.4	75	12	55	26	32
MeA	Metamora-Capac sandy loams, 0 to 4 percent slopes	8.40	3.3%		1.5ft.	Somewhat poorly drained	Ile	4.8	120	19	100	36	60
Sb	Sebewa loam, 0 to 2 percent slopes	7.97	3.1%		0ft.	Poorly drained	IIw						
Gf	Gilford sandy loam, 0 to 2 percent slopes, gravelly subsoil	7.54	2.9%		3.5ft.	Poorly drained	IIlw		149				42 59
CaA	Capac loam, 0 to 4 percent slopes	4.48	1.7%		1ft.	Somewhat poorly drained	Ile						
Ho	Houghton muck, 0 to 1 percent slopes	3.58	1.4%		0ft.	Very poorly drained	Vw						
W	Water	2.85	1.1%		> 6.5ft.								
Weighted Average								1.7	41.7	6.3	28.8	14.4	18.4

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.