



# Soils Map



State: **Indiana**  
 County: **Clinton**  
 Location: **22-23N-1W**  
 Township: **Owen**  
 Acres: **1037.64**  
 Date: **2/23/2023**



Soils data provided by USDA and NRCS.

Area Symbol: IN015, Soil Area Version: 27											
Area Symbol: IN023, Soil Area Version: 26											
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Non-Irr Class *c	Corn Bu	Soybeans Bu	Winter wheat Bu	*n NCCPI Corn	*n NCCPI Soybeans
FdA	Fincastle-Crosby silt loams, 0 to 2 percent slopes	184.76	17.8%		0.5ft.	llw	156	51	69	81	70
MtB	Miami-Crosby silt loams, 2 to 6 percent slopes	176.76	17.0%		2.5ft.	lle	147	50	65	70	59
Mp	Milford silty clay loam, occasionally flooded	109.71	10.6%		0.2ft.	llw	140	37	56	98	76
MdB2	Martinsville, till substratum-Miami loams, 2 to 6 percent slopes, eroded	91.82	8.8%		2.7ft.	lle	135	47	65	79	60
Ty	Treaty silt loam, 0 to 2 percent slopes	85.05	8.2%		0ft.	llw	177	62	61	78	72
CwB	Crosby-Fincastle silt loams, 1 to 3 percent slopes	61.65	5.9%		1.2ft.	lle	140	46	63	76	59
So	Sloan silt loam, occasionally flooded	56.07	5.4%		0ft.	lllw	155	42	62	54	80
Cz	Cyclone silty clay loam, 0 to 2 percent slopes	48.64	4.7%		0.3ft.	llw	185	65	75	87	81
MfC3	Martinsville, till substratum-Miami clay loams, 6 to 12 percent slopes, severely eroded	32.87	3.2%		2.7ft.	lve	121	43	58	68	46
CbA	Camden variant silt loam, 0 to 2 percent slopes	26.70	2.6%		> 6.5ft.	l	155	54	78	90	79
Ce	Ceresco loam	24.81	2.4%		1.2ft.	lllw	113	30	51	58	67
Wh	Whitaker silt loam, 0 to 2 percent slopes	22.29	2.1%		0.5ft.	llw	155	51	70	92	78

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Non-Irr Class *c	Corn Bu	Soybeans Bu	Winter wheat Bu	*n NCCPI Corn	*n NCCPI Soybeans
RrB2	Rockfield-Williamstown complex, 1 to 6 percent slopes, eroded	12.88	1.2%		2.2ft.	lle	133	47	60	72	54
Cg	Ceresco fine sandy loam, occasionally flooded	12.35	1.2%		1.2ft.	llw	108	28	49	57	66
Mx	Milford silty clay loam	11.09	1.1%		0.2ft.	llw	160	44	64	83	75
Mo	Milford silt loam, pothole	10.65	1.0%		0.2ft.	llw	80	22	32	75	35
Ck	Ceresco variant fine sandy loam, occasionally flooded	9.08	0.9%		1.2ft.	llw	111	29	50	53	59
Lo	Landes loam, moderately wet, occasionally flooded	9.04	0.9%		5ft.	llw	106	34	53	69	57
XeB	Xenia silt loam, 2 to 4 percent slopes	8.40	0.8%		1.8ft.	lle	156	54	69	83	73
Ph	Patton silty clay loam, 0 to 2 percent slopes	6.44	0.6%		0.5ft.	llw	173	51	71	76	73
MwA	Miami-Martinsville silt loams, 0 to 2 percent slopes	6.11	0.6%		> 6.5ft.	l	127	44	64	82	74
RuB	Russell silt loam, 2 to 6 percent slopes	5.52	0.5%		4.7ft.	lle	156	54	69	87	76
FsC	Fox loam, 6 to 15 percent slopes	4.47	0.4%		> 6.5ft.	llle	75	26	38	63	56
FaA	Fincastle-Starks silt loams, 0 to 2 percent slopes	4.42	0.4%		0.5ft.	llw	166	54	74	90	78
MnC	Miami silt loam, 6 to 12 percent slopes	4.25	0.4%		2.7ft.	llle	138	48	61	63	57
FtC3	Fox gravelly clay loam, 6 to 15 percent slopes, severely eroded	4.18	0.4%		> 6.5ft.	llle	73	25	37	54	38
RoA	Rockfield silt loam, 0 to 2 percent slopes	3.31	0.3%		2.2ft.	l	145	51	66	87	72
Wk	Washtenaw silt loam	2.22	0.2%		0.2ft.	llw	165	49	66	76	61
Cp	Cohoctah loam, gravelly substratum, occasionally flooded	1.09	0.1%		0.2ft.	llw	155	33	62	66	62
Wa	Walkill silt loam	0.88	0.1%		0.2ft.	lllw	140	40		90	78
Ma	Mahalasville silty clay loam, 0 to 2 percent slopes	0.13	0.0%		0.3ft.	llw	175	61	58	85	77
<b>Weighted Average</b>						<b>2.12</b>	<b>147.6</b>	<b>47.8</b>	<b>63.5</b>	<b>*n 77.5</b>	<b>*n 67.1</b>

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method