



Soil Map



State: **Michigan**
 County: **Shiawassee**
 Location: **4-6N-2E**
 Township: **Bennington**
 Acres: **388.07**
 Date: **5/26/2015**

Maps Provided By:



Soils data provided by USDA and NRCS.

Viewing soils data as of 1/21/2012

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
ChB	Celina loam, 2 to 6 percent slopes	90.54	23.3%		4.5ft.	Ile	4.8	132	19	95	35	66
CtA	Conover loam, 0 to 2 percent slopes	59.79	15.4%		1ft.	IIlw	5	143	19	105	40	71
CtB	Conover loam, 2 to 6 percent slopes	44.65	11.5%		1ft.	Ile	5	143	19	105	40	71
Gg	Gilford sandy loam	42.24	10.9%		0ft.	IIIw	4.8	132	19	100	35	61
BmA	Boyer loamy sand, 0 to 2 percent slopes	42.19	10.9%		> 6.5ft.	IIIs	3.8	88	14	60	30	38
WeA	Wasepi sandy loam, 0 to 2 percent slopes	29.83	7.7%		1ft.	IIIw	4.5	126	19	100	35	60
Sd	Sebewa loam	26.24	6.8%		0ft.	IIlw	4.8	132	19	100	35	61
Bw	Brookston loam	19.09	4.9%		0ft.	IIlw	5.5	154	22	115	45	71
BmB	Boyer loamy sand, 2 to 6 percent slopes	12.63	3.3%		> 6.5ft.	IIIe	3.8	88	14	60	30	38
MuC2	Miami loam, 6 to 12 percent slopes, moderately eroded	12.36	3.2%		> 6.5ft.	IIIe	4.6	125	18	90	33	63
KhB	Kendallville sandy loam, 2 to 6 percent slopes	3.49	0.9%		> 6.5ft.	Ile	3.5	126	19	95	35	60
BrC	Boyer sandy loam, 6 to 12 percent slopes	2.03	0.5%		> 6.5ft.	IIIe	3.4	82	12	55	26	35
KhC2	Kendallville sandy loam, 6 to 12 percent slopes, moderately eroded	1.94	0.5%		> 6.5ft.	IIIe	3.2	115	17	86	32	54
BmC	Boyer loamy sand, 6 to 12 percent slopes	0.96	0.2%		> 6.5ft.	IIIe	3.4	82	12	55	26	35
MuB2	Miami loam, 2 to 6 percent slopes, moderately eroded	0.09	0.0%		> 6.5ft.	Ile	4.6	125	18	90	33	63
Weighted Average							4.7	128.6	18.3	94.5	36	61.8

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.

Soils data provided by USDA and NRCS.