



# Soils Map



State: **Indiana**  
 County: **Warren**  
 Location: **34-23N-8W**  
 Township: **Pine**  
 Acres: **191.89**  
 Date: **7/20/2017**



Soils data provided by USDA and NRCS.

©2017 AgriData, Inc.

## Area Symbol: IN171, Soil Area Version: 19

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
RfB2	Rainsville-Williamstown-Rockfield silt loams, 2 to 6 percent slopes, eroded	68.11	35.5%		2.2ft.	Moderately well drained	Ile	133	47	60
SIA	Starks silt loam, till substratum, 0 to 2 percent slopes	26.28	13.7%		1.2ft.	Somewhat poorly drained	Ilw	160	52	72
WtC2	Williamstown-Rainsville silt loams, 6 to 12 percent slopes, eroded	19.51	10.2%		2ft.	Moderately well drained	Ille	122	43	55
MdA	Martinsville loam, 0 to 2 percent slopes	14.69	7.7%		> 6.5ft.	Well drained	I	140	49	71
MpC3	Miami clay loam, 6 to 12 percent slopes, severely eroded	11.21	5.8%		2.5ft.	Moderately well drained	Ive	121	41	54
Cz	Cyclone silty clay loam, 0 to 2 percent slopes	8.47	4.4%		0.2ft.	Poorly drained	Ilw	184	54	75
Mr	Milford silty clay loam, pothole	7.91	4.1%		0.2ft.	Very poorly drained	Ilw	80	22	32
Hm	Houghton muck, drained	7.38	3.8%		0ft.	Very poorly drained	Illw	155	42	62
RoB2	Rockfield silt loam, 2 to 6 percent slopes, eroded	4.77	2.5%		2.2ft.	Moderately well drained	Ile	141	50	64
WIA	Waupecan silt loam, moderately wet, 0 to 2 percent slopes	3.70	1.9%		3.2ft.	Moderately well drained	I	160	52	72
RdA	Rainsville silt loam, 0 to 2 percent slopes	3.33	1.7%		2.2ft.	Moderately well drained	I	140	49	63
OcB2	Ockley silt loam, 2 to 6 percent slopes, eroded	3.25	1.7%		> 6.5ft.	Well drained	Ile	132	46	65
Dy	Du Page loam, frequently flooded	3.16	1.6%		> 6.5ft.	Well drained	Ilw	131	41	
CbB2	Camden silt loam, 2 to 6 percent slopes, eroded	3.06	1.6%		> 6.5ft.	Well drained	Ile	141	50	70
Wh	Washtenaw silt loam	2.40	1.3%		0.2ft.	Poorly drained	Ilw	165	49	66
MoE2	Miami loam, 15 to 25 percent slopes, eroded	1.55	0.8%		> 6.5ft.	Well drained	Vle			
Pm	Peotone silty clay loam, pothole	1.12	0.6%		0.2ft.	Very poorly drained	Illw	85	11	34
OcA	Ockley silt loam, 0 to 2 percent slopes	0.77	0.4%		> 6.5ft.	Well drained	Ie	134	47	66
HeG	Hennepin loam, 30 to 70 percent slopes	0.58	0.3%		> 6.5ft.	Well drained	Vlle			



Cs	Comfrey loam, stratified substratum, frequently flooded, undrained	0.20	0.1%		0.2ft.	Very poorly drained	Vw			
We	Wallkill variant silty clay loam	0.15	0.1%		0.2ft.	Very poorly drained	Illw	160	43	64
RpG	Rodman gravelly loam, 25 to 60 percent slopes	0.12	0.1%		> 6.5ft.	Excessively drained	Vllc			
Be	Beaucoup silty clay loam, frequently flooded, undrained	0.10	0.1%		0.2ft.	Very poorly drained	Vw			
Bk	Beckville loam, occasionally flooded	0.07	0.0%		2.2ft.	Moderately well drained	Ilw			
<b>Weighted Average</b>								<b>135.6</b>	<b>45.5</b>	<b>60</b>

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.