



Soils Map



Soils data provided by USDA and NRCS.



State: **Indiana**
 County: **Warren**
 Location: **34-23N-8W**
 Township: **Pine**
 Acres: **186.71**
 Date: **1/13/2023**



Maps Provided By:



Archived Soils Ending 1/21/2012

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn Bu	Soybeans Bu	Winter wheat Bu
RfB2	Rainsville-Williamstown-Rockfield silt loams, 2 to 6 percent slopes, eroded	68.39	36.6%		2.2ft.	4.6ft. (Densic material)	Moderately well drained	Ile	142	50	64
SIA	Starks silt loam, till substratum, 0 to 2 percent slopes	25.89	13.9%		1.2ft.	4.6ft. (Densic material)	Somewhat poorly drained	IIw	165	54	74
WtC2	Williamstown-Rainsville silt loams, 6 to 12 percent slopes, eroded	19.08	10.2%		2ft.	2.9ft. (Densic material)	Moderately well drained	IIIe	130	46	59
MdA	Martinsville loam, 0 to 2 percent slopes	14.44	7.7%		> 6.5ft.	> 6.5ft.	Well drained	I	145	51	73
MpC3	Miami clay loam, 6 to 12 percent slopes, severely eroded	9.30	5.0%		2.7ft.	2.1ft. (Densic material)	Moderately well drained	IVe	125	44	56
Cz	Cyclone silty clay loam	8.43	4.5%		0.5ft.	> 6.5ft.	Poorly drained	IIw	190	54	76
Mr	Milford silty clay loam, pothole	7.89	4.2%		0.2ft.	> 6.5ft.	Very poorly drained	IIw	160	44	64
Hm	Houghton muck, drained	5.42	2.9%		0ft.	> 6.5ft.	Very poorly drained	IIIw	155	42	62
RoB2	Rockfield silt loam, 2 to 6 percent slopes, eroded	4.55	2.4%		2.2ft.	4.6ft. (Densic material)	Moderately well drained	Ile	150	53	68

Soils data provided by USDA and NRCS.

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn Bu	Soybeans Bu	Winter wheat Bu
WIA	Waupecan silt loam, moderately wet, 0 to 2 percent slopes	3.73	2.0%		3.2ft.	5.1ft. (Strongly contrasting textural stratification)	Moderately well drained	I	165	54	74
RdA	Rainsville silt loam, 0 to 2 percent slopes	3.33	1.8%		2.2ft.	4.6ft. (Densic material)	Moderately well drained	I	145	51	65
OcB2	Ockley silt loam, 2 to 6 percent slopes, eroded	3.28	1.8%		> 6.5ft.	4.9ft. (Strongly contrasting textural stratification)	Well drained	Ile	125	44	63
Dy	Du Page loam, frequently flooded	3.02	1.6%		> 6.5ft.	> 6.5ft.	Well drained	IIw	135	42	
CbB2	Camden silt loam, 2 to 6 percent slopes, eroded	2.95	1.6%		> 6.5ft.	> 6.5ft.	Well drained	Ile	150	53	75
Wh	Washtenaw silt loam	2.41	1.3%		0.2ft.	> 6.5ft.	Poorly drained	IIw	165	49	66
MoE2	Miami loam, 15 to 25 percent slopes, eroded	1.57	0.8%		> 6.5ft.	2.4ft. (Densic material)	Well drained	VIle			
Pm	Peotone silty clay loam, pothole	1.12	0.6%		0.2ft.	> 6.5ft.	Very poorly drained	IIIw	165	44	66
OcA	Ockley silt loam, 0 to 2 percent slopes	0.77	0.4%		> 6.5ft.	4ft. (Strongly contrasting textural stratification)	Well drained	I	136	47	68
HeG	Hennepin loam, 30 to 70 percent slopes	0.61	0.3%		> 6.5ft.	> 6.5ft.	Well drained	VIIle			
Cs	Comfrey loam, stratified substratum, frequently flooded, undrained	0.22	0.1%		0.2ft.	> 6.5ft.	Very poorly drained	V w			
RpG	Rodman gravelly loam, 25 to 60 percent slopes	0.15	0.1%		> 6.5ft.	1.2ft. (Strongly contrasting textural stratification)	Excessively drained	VIIle			
Be	Beaucoup silty clay loam, frequently flooded, undrained	0.10	0.1%		0.2ft.	> 6.5ft.	Very poorly drained	V w			
Bk	Beckville loam, occasionally flooded	0.06	0.0%		2.2ft.	> 6.5ft.	Moderately well drained	IIw			
Weighted Average								2.18	145.4	48.8	64.2

*c: Using Capabilities Class Dominant Condition Aggregation Method