
Soils Map





State: Ohio

County: Montgomery

8-3N-4E Location: Township: German Acres: 39.74 6/28/2016 Date:





Soils data provided by USDA and NRCS.

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Corn	Oats	Soybeans	Winter wheat	*eFOTG PI
CeB	Celina silt loam, 2 to 6 percent slopes	13.03	32.8%		2ft.	Moderately well drained	lle	131	70	46	58	0
MIB2	Miamian silt loam, 2 to 6 percent slopes, eroded	10.56	26.6%		3.1ft.	Well drained	lle	119		36	48	0
Вр	Brookston silt loam	7.57	19.0%		Oft.	Very poorly drained	llw	115	80	40	45	92
MIB	Miamian silt loam, 2 to 6 percent slopes	3.05	7.7%		2.5ft.	Well drained	lle	130		46	59	0
CeA	Celina silt loam, 0 to 2 percent slopes	2.42	6.1%		2.2ft.	Moderately well drained	I	115	80	40	45	79
MnD3	Miamian clay loam, 12 to 18 percent slopes, severely eroded	0.99	2.5%		> 6.5ft.	Well drained	Vle		45			47
CsA	Crosby silt loam, Southern Ohio Till Plain, 0 to 2 percent slopes	0.79	2.0%		1.5ft.	Somewhat poorly drained	llw	120		46	48	0
MnC3	Miamian clay loam, 6 to 12 percent slopes, severely eroded	0.77	1.9%		3.1ft.	Well drained	IVe	100		35	45	0
Bs	Brookston silty clay loam, fine texture, 0 to 2 percent slopes	0.44	1.1%		4.5ft.	Poorly drained	llw	129		48	51	0
HmF3	Hennepin and Miamian soils, 18 to 50 percent slopes, severely eroded	0.12	0.3%		> 6.5ft.	Well drained	VIIe					0
Weighted Average									44.2	40.4	50	23.5


Area Symbol: OH113, Soil Area Version: 14


*eftog PI: Obtained from the NRCS eFOTG (http://efotg.sc.egov.usda.gov)

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.

Soil Map





State: Ohio
County: Warren
Location: 29-2N-5E
Township: Franklin
Acres: 44.77
Date: 3/20/2015





Soils data provided by USDA and NRCS.

	. ,										
Area Symbol: OH113, Soil Area Version: 13 Area Symbol: OH165, Soil Area Version: 13											
Code	Soil Description	Acres	Percent of field	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class	Corn	Soybeans	Winter wheat	*eFOTG PI
Rn	Ross loam	18.92	42.3%	5ft.	> 6.5ft.	Well drained	llw	120	40	40	91
AbA	Abscota sand, calcareous variant	11.01	24.6%	4.5ft.	> 6.5ft.	Well drained	IVs	70	25	35	51
Gd	Genesee fine sandy loam	5.54	12.4%	> 6.5ft.	> 6.5ft.	Well drained	llw	115	40	40	67
Rs	Ross silt loam	4.41	9.9%	> 6.5ft.	> 6.5ft.	Well drained	llw	125	40	45	92
WaA	Warsaw loam, 0 to 2 percent slopes	2.05	4.6%	> 6.5ft.	3ft. (Strongly contrasting textural stratification)	Well drained	lls	110	40	40	74
Ld	Landes sandy loam	1.98	4.4%	> 6.5ft.	> 6.5ft.	Well drained	llw	103	34	39	67
FIB	Fox loam, 2 to 6 percent slopes	0.48	1.1%	> 6.5ft.	2.4ft. (Strongly contrasting textural stratification)	Well drained	lle	90	30	40	69
WaA	Warsaw silt loam, 0 to 2 percent slopes	0.38	0.8%	> 6.5ft.	2.6ft. (Strongly contrasting textural stratification)	Well drained	lls	110	40	40	74
Weighted Average									35.9	39.2	76.1

Area Symbol: OH113, Soil Area Version: 13 Area Symbol: OH165, Soil Area Version: 13

*eftog PI: Obtained from the NRCS eFOTG (http://efotg.sc.egov.usda.gov)

Soils data provided by USDA and NRCS.