



Soils Map



State: **Indiana**
 County: **Miami**
 Location: **15-28N-4E**
 Township: **Richland**
 Acres: **309.63**
 Date: **2/12/2017**



Soils data provided by USDA and NRCS.

Area Symbol: IN103, Soil Area Version: 20

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Soil Drainage | Non-Irr Class *c | Corn | Soybeans | Winter wheat | Oats | Corn silage |
|-------------------------|--|--------|------------------|----------------------|-------------|-------------------------|------------------|--------------|-------------|--------------|------------|-------------|
| Re | Rensselaer loam | 138.84 | 44.8% | | 0.7ft. | Poorly drained | IIw | 172 | 48 | 69 | | |
| OtA | Oshtemo sandy loam, 0 to 4 percent slopes | 80.40 | 26.0% | | > 6.5ft. | Well drained | IIIs | 99 | 35 | 50 | | |
| MtC3 | Morley silty clay loam, 6 to 12 percent slopes, severely eroded | 30.19 | 9.8% | | 2.5ft. | Moderately well drained | IVe | 105 | 37 | 47 | | |
| MeB | Metea loamy fine sand, 2 to 6 percent slopes | 15.68 | 5.1% | | > 6.5ft. | Well drained | IIIe | 110 | 39 | 55 | | |
| MsB | Glynnwood silt loam, 2 to 6 percent slopes | 11.71 | 3.8% | | 1.5ft. | Moderately well drained | Ile | 133 | 46 | 60 | | |
| Sn | Sleeth loam | 6.93 | 2.2% | | 1.2ft. | Somewhat poorly drained | IIw | 140 | 46 | 63 | | |
| Pw | Pewamo silty clay loam, 0 to 1 percent slopes | 4.02 | 1.3% | | 0.5ft. | Very poorly drained | IIw | 157 | 47 | 64 | | |
| Pm | Palms muck, drained | 3.95 | 1.3% | | 0.7ft. | Very poorly drained | IIIw | 158 | 43 | 63 | | |
| OsB | Ormas-Oshtemo loamy sands, 2 to 8 percent slopes | 3.74 | 1.2% | | > 6.5ft. | Well drained | IIIs | 82 | 29 | 41 | | |
| So | Sloan silty clay loam | 3.59 | 1.2% | | 0ft. | Very poorly drained | IIIw | 150 | 40 | | | |
| MtD3 | Morley silty clay loam, 12 to 25 percent slopes, severely eroded | 3.40 | 1.1% | | 2.5ft. | Moderately well drained | VIe | 75 | 26 | 34 | | |
| Au | Aubbeenaubbee sandy loam, 0 to 2 percent slopes | 2.39 | 0.8% | | 1.2ft. | Somewhat poorly drained | IIw | 135 | 44 | 61 | | |
| MrB | Morley sandy loam, 2 to 6 percent slopes | 2.38 | 0.8% | | 2.7ft. | Moderately well drained | Ile | 117 | 41 | 53 | | |
| Ba | Blount loam, 1 to 3 percent slopes | 1.38 | 0.4% | | 2ft. | Somewhat poorly drained | Ile | 140 | 46 | 63 | 61 | 1 |
| Wh | Washtenaw silt loam | 1.03 | 0.3% | | 0.2ft. | Poorly drained | IIw | 165 | 49 | 66 | | |
| Weighted Average | | | | | | | | 137.5 | 42.2 | 58.9 | 0.3 | *- |

Area Symbol: IN103, Soil Area Version: 20

*c: Using Capabilities Class Dominant Condition Aggregation Method