

Soils Map





State: Ohio
County: Williams
Location: 5-10S-1W
Township: Mill Creek
Acres: 292.16
Date: 11/16/2016







Area Symbol: OH171, Soil Area Version: 12												
	Soil Description		Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat	*eFOTG PI
BnB	Blount loam, 2 to 6 percent slopes	63.82	21.8%		0.7ft.	2.9ft. (Densic material)	Somewhat poorly drained	lle	140	46	63	74
Mh	Millgrove loam	39.76	13.6%		0.2ft.	> 6.5ft.	Very poorly drained	llw	125	40	50	100
Pm	Pewamo silty clay loam, 0 to 1 percent slopes	38.34	13.1%		0.5ft.	> 6.5ft.	Very poorly drained	llw	157	47	64	83
BnA	Blount loam, 0 to 2 percent slopes	23.73	8.1%		0.7ft.	3.1ft. (Densic material)	Somewhat poorly drained	llw	141	46	63	77
DgA	Digby sandy loam, 0 to 3 percent slopes	23.00	7.9%		1ft.	> 6.5ft.	Somewhat poorly drained	llw	110	40	46	75
Gf	Gilford fine sandy loam, till plain, 0 to 2 percent slopes	22.53	7.7%		Oft.	> 6.5ft.	Poorly drained	IIIw	146	33	59	83
HkA	Haskins sandy loam, 0 to 3 percent slopes	18.67	6.4%		1.2ft.	> 6.5ft.	Somewhat poorly drained	llw	105	40	46	73
GIB	Glynwood loam, 2 to 6 percent slopes	16.29	5.6%		1.5ft.	2.9ft. (Densic material)	Moderately well drained	lle	133	43	57	69
OtB	Ottokee fine sand, 0 to 6 percent slopes	10.97	3.8%		2ft.	> 6.5ft.	Moderately well drained	IIIs	90	34	40	63
Md	Mermill loam	10.90	3.7%		0.2ft.	> 6.5ft.	Very poorly drained	llw	125	44	50	92
GIC2	Glynwood loam, 6 to 12 percent slopes, moderately eroded	8.13	2.8%		2.2ft.	> 6.5ft.	Moderately well drained	IVe	85	28	32	61
НаВ	Haney loam, 1 to 6 percent slopes	4.10	1.4%		2.2ft.	> 6.5ft.	Moderately well drained	lle	100	40	48	71
HnA	Haskins loam, 0 to 3 percent slopes	3.85	1.3%		1.2ft.	> 6.5ft.	Somewhat poorly drained	llw	110	42	46	76
GIC	Glynwood loam, 6 to 12 percent slopes	2.13	0.7%		2.2ft.	> 6.5ft.	Moderately well drained	IIIe	90	33	40	64



То	Toledo silty clay, 0 to 1 percent slopes	1.96	0.7%		1.7ft.	> 6.5ft.	Very poorly drained	IIIw	117	39.9	45	75
SdB	Seward loamy fine sand, 2 to 6 percent slopes	1.49	0.5%		2ft.	> 6.5ft.	Moderately well drained	lle	90	30	34	59
GIB2	Glynwood loam, 2 to 6 percent slopes, eroded	0.99	0.3%		1.5ft.	2.5ft. (Densic material)	Moderately well drained	lle	128	41	55	67
RIB	Rawson sandy loam, 2 to 6 percent slopes	0.96	0.3%		2.7ft.	> 6.5ft.	Moderately well drained	lle	100	35	42	70
Ca	Carlisle muck, disintegration moraine, drained, 0 to 2 percent slopes	0.54	0.2%		Oft.	> 6.5ft.	Very poorly drained	IIIw	127	44		66
Weighted Average									129.9	41.9	55	79.1

Area Symbol: OH171, Soil Area Version: 12

*eftog PI: Obtained from the NRCS eFOTG (http://efotg.sc.egov.usda.gov)

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.