

State: Wisconsin County: Fond Du Lac Location: 23-15N-14E Township: Metomen Acres: 68.85 8/28/2017



Date:







Soils data provided by USDA and NRCS.

Area	Symbol: WI039, Soil Area	Versic	n: 16								
	Soil Description		Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
LvB	Lomira silt loam, 2 to 6 percent slopes	24.18	35.1%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
VgA	Virgil silt loam, 0 to 2 percent slopes	7.92	11.5%		2ft.	> 6.5ft.	Somewhat poorly drained		155	51	48
ScB	St. Charles silt loam, 2 to 6 percent slopes	7.67	11.1%		4ft.	> 6.5ft.	Well drained	lle	145	48	51
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	7.26	10.5%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
LvC2	Lomira silt loam, 6 to 12 percent slopes, eroded	6.81	9.9%		> 6.5ft.	> 6.5ft.	Well drained	IIIe	106.2	35.2	41.5
PhA	Pella silt loam, 0 to 2 percent slopes	6.57	9.5%		Oft.	> 6.5ft.	Poorly drained	llw	124.4	46	
BsA	Brookston silt loam, 0 to 2 percent slopes	3.13	4.5%		0.7ft.	> 6.5ft.	Poorly drained	llw	120	40	
ScA	St. Charles silt loam, 0 to 2 percent slopes	2.80	4.1%		4ft.	> 6.5ft.	Well drained	lw	150	48	51
JuA	Juneau silt loam, 0 to 2 percent slopes	2.51	3.6%		Oft.	> 6.5ft.	Moderately well drained	1	145	48	50
	•	•	•			•	Weight	ed Average	125.7	42	39.8

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin
County: Fond Du Lac
Location: 36-16N-16E
Township: Eldorado
Acres: 33.99
Date: 8/22/2017









Soils data provided by USDA and NRCS.

Area	Symbol: WI039, Soil Are	a Vers	sion: 16										
Code	•	Acres		Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
KoB2	Kewaunee silty clay loam, 2 to 6 percent slopes, eroded	15.40	45.3%		5.4ft.	Well drained	lle	4.8	108.8	18	68.8	35.8	41.5
МсВ	Manawa silty clay loam, 2 to 6 percent slopes	9.38	27.6%		1ft.	Somewhat poorly drained	lle	4.5	115	19	65	38	39
KoA	Kewaunee silty clay loam, 0 to 2 percent slopes	4.06	11.9%		5.8ft.	Well drained	lls	4.8	110	18	70	36	42
KnB	Kewaunee silt loam, 2 to 6 percent slopes	2.62	7.7%		5.4ft.	Well drained	lle	4.8	110	18	70	36	42
KoB	Kewaunee silty clay loam, 2 to 6 percent slopes	2.53	7.4%		5.8ft.	Well drained	lle	4.8	110	18	70	36	42
						Weighte	d Average	4.7	110.8	18.3	68.1	36.5	40.9

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin County: Fond Du Lac Location: 18-14N-16E Township: **Oakfield** Acres: 72.18 8/22/2017 Date:








Soils data provided by USDA and NRCS.

Alea S	<u>ymbol: WI039, Soil Are</u>	<del>ç</del> a ver	<u> 51011. 10</u>		,								
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
EsB	Elliott silt loam, 2 to 6 percent slopes	21.21	29.4%		2ft.	Somewhat poorly drained	1	4.7	140	23	75	46	45
VgA	Virgil silt loam, 0 to 2 percent slopes	13.45	18.6%		2ft.	Somewhat poorly drained		5.4	155	25	80	51	48
AtA	Ashkum silty clay loam, 0 to 3 percent slopes	8.22	11.4%		0.5ft.	Poorly drained	llw		130	21		43	
ScA	St. Charles silt loam, 0 to 2 percent slopes	8.14	11.3%		4ft.	Well drained	lw	5.5	150	25	85	48	51
EsA	Elliott silt loam, 0 to 2 percent slopes	6.50	9.0%		2ft.	Somewhat poorly drained		4.7	140	23	75	46	45
PhA	Pella silt loam, 0 to 2 percent slopes	5.24	7.3%		Oft.	Poorly drained	llw		124.4	20		46	
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	5.00	6.9%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45
ВсВ	Beecher silt loam, 2 to 6 percent slopes	1.71	2.4%		2ft.	Somewhat poorly drained		4.7	135	22	75	45	45
MoB	Mayville silt loam, 2 to 6 percent slopes	1.10	1.5%		2.1ft.	Moderately well drained	lle	5	131	21	78	43	49
MzdB2	Morley silt loam, 2 to 6 percent slopes, eroded	1.01	1.4%		3.3ft.	Moderately well drained	lle	4.7	125	20	75	41	45
PnA	Pella silty clay loam, cool, 0 to 2 percent slopes	0.46	0.6%		Oft.	Poorly drained	llw		162	25	87	53.5	66.5
LvB	Lomira silt loam, 2 to 6 percent slopes	0.14	0.2%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method



25 30 29 ∘s -Metovale-Rd-Тс 32 <sup>6</sup>©2017 AgriData, Iກົc.

State: Wisconsin County: Fond Du Lac Location: 31-15N-15E Township: **Springvale** 

Acres: 53.29 8/22/2017 Date:








Area	Symbol: WI039, Soil Are	a Vers	sion: 16										
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
LtC3	LeRoy soils, 6 to 12 percent slopes, severely eroded	15.98	30.0%		5.8ft.	Well drained	IVe	3.2	80	13	55	26	36
VgA	Virgil silt loam, 0 to 2 percent slopes	9.37	17.6%		2ft.	Somewhat poorly drained	llw	5.4	155	25	80	51	48
LrB2	LeRoy silt loam, 2 to 6 percent slopes, eroded	9.34	17.5%		5.8ft.	Well drained	lle	3.8	95	15	70	33	42
PhA	Pella silt loam, 0 to 2 percent slopes	7.97	15.0%		Oft.	Poorly drained	llw		124.4	20		46	
LvB	Lomira silt loam, 2 to 6 percent slopes	4.82	9.0%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	2.48	4.7%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45
PsA	Plano silt loam, till substratum, 0 to 2 percent slopes	1.78	3.3%		3.8ft.	Well drained	I						
LtD3	LeRoy soils, 12 to 20 percent slopes, severely eroded	1.32	2.5%		5.8ft.	Well drained	Vle	2.8	70	12	45	23	32
HmD	Hochheim loam, 12 to 20 percent slopes	0.23	0.4%		> 6.5ft.	Well drained	IVe						
						Weighte	d Average	3.2	104	16.8	54.2	35.2	33.6

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method



2.8%

2.2%

1.1%

1.0%

0.9%

0.9%

6.32

5.04

2.53

2.19

2.06

2.04

Area Symbol: WI039, Soil Area Version: 16

St. Charles silt loam, 2 to 6

Washtenaw silt loam, 0 to 2

Casco loam, 2 to 6 percent

Lomira silt loam, 12 to 20

percent slopes, severely

Lomira silt loam, 6 to 12

percent slopes, eroded
Theresa-Casco silt loams, 12

to 20 percent slopes, eroded

percent slopes

percent slopes

slopes


ScB

WsA

CeB

LvD3

LvC2



State: Wisconsin
County: Fond Du Lac
Location: 7-15N-15E
Township: Springvale
Acres: 229.29
Date: 8/28/2017



Well drained

Poorly drained

excessively drained

Somewhat

Well drained

Well drained

Well drained

> 6.5ft.

> 6.5ft.

> 6.5ft.

> 6.5ft.

> 6.5ft.

1.4ft. (Strongly contrasting

textural stratification)

145

130

90

90

95

lle

llw

Ille

IVe

IIIe 106.2

IVe

48

43

30

30

35.2

32

51

36

35

41.5

36





Non-Irr Water Soil Drainage Winter Soil Description Acres Percent of Non-Irr Restrictive Layer Corn Soybeans Code field Class Table Class \*c wheat Legend LvB Lomira silt loam, 2 to 6 53.84 23.5% > 6.5ft > 6.5ft. Well drained lle 115 38 45 percent slopes VgA Virgil silt loam, 0 to 2 percent 40.45 17.6% 2ft. > 6.5ft. Somewhat poorly 155 51 48 llw slopes drained St. Charles silt loam, 0 to 2 ScA 29.01 12.7% 4ft. > 6.5ft. Well drained lw 150 48 51 percent slopes Lomira silt loam, 2 to 6 27.37 11.9% > 6.5ft. > 6.5ft Well drained 115 38 45 LvB2 lle percent slopes, eroded Pella silt loam, 0 to 2 percent 11.1% PhA 25.56 Oft. > 6.5ft. Poorly drained 124.4 46 llw Theresa-Casco silt loams, 6 18.01 7.9% 5.8ft. Ille 105 35 40 TsC2 > 6.5ft. Well drained to 12 percent slopes, eroded Lomira silt loam, 6 to 12 33 39 LvC3 12.16 5.3% > 6.5ft > 6.5ft. Well drained IVe 100 percent slopes, severely eroded

4ft.

Oft.

> 6.5ft.

> 6.5ft


> 6.5ft.


5.8ft.



Hu	Houghton mucky peat, 0 to 2 percent slopes	1.43	0.6%		Oft.	> 6.5ft.	Very poorly drained	IIIw	135	46	
TsB2	Theresa-Casco silt loams, 2 to 6 percent slopes, eroded	0.98	0.4%		5.8ft.	> 6.5ft.	Well drained	lle	115	38	44
Rw	Rollin mucky peat	0.12	0.1%		Oft.	> 6.5ft.	Very poorly drained	IVw	100		
JuA	Juneau silt loam, 0 to 2 percent slopes	0.09	0.0%		Oft.	> 6.5ft.	Moderately well drained	I	145	48	50
		0.09	0.0%		5.8ft.	> 6.5ft.	Excessively drained	IVs			
RoD2 Rodman gravelly loam, 12 to 20 percent slopes, eroded 0.09 0.0% 5.8ft. > 6.5ft. Excessively drained IVs										42.1	39.1

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin
County: Fond Du Lac
Location: 19-15N-15E
Township: Springvale
Acres: 454.02
Date: 8/28/2017









Area Symbol: WI039, Soil Area Version: 16 Non-Irr Class Soil Drainage Percent of Water Winter Code Soil Description Restrictive Non-Irr Corn Soybeans field Legend Table Layer Class \*c wheat LvB Lomira silt loam, 2 to 6 percent 126.09 27.8% > 6.5ft. > 6.5ft 38 45 Well drained lle 115 51 VgA Virgil silt loam, 0 to 2 percent 102.55 22.6% 2ft. > 6.5ft Somewhat poorly llw 155 48 LvB2 Lomira silt loam, 2 to 6 percent 68.53 15.1% > 6.5ft. > 6.5ft Well drained lle 115 38 45 slopes, eroded PhA Pella silt loam, 0 to 2 percent 54.60 12.0% Oft. > 6.5ft Poorly drained llw 124.4 46 slopes ScA St. Charles silt loam, 0 to 2 26.73 5.9% 4ft. > 6.5ft Well drained lw 150 48 51 percent slopes Lomira silt loam, 6 to 12 percent 41.5 Ille 106.2 35.2 LvC2 22.87 5.0% > 6.5ft. > 6.5ft Well drained slopes, eroded Lomira silt loam, 0 to 2 percent 120 40 47 I vA 22.69 5.0% > 6.5ft. > 6.5ft Well drained lw LeRoy soils, 6 to 12 percent 80 26 LtC3 6.53 1.4% 5.8ft. > 6.5ft Well drained IVe 36 slopes, severely eroded ScB St. Charles silt loam, 2 to 6 5.90 1.3% 4ft. > 6.5ft Well drained lle 145 48 51 percent slopes Virgil silt loam, 2 to 6 percent 5.34 1.2% 2ft. > 6.5ft 155 51 48 VgB Somewhat poorly llw slopes drained Lomira silt loam, 6 to 12 percent > 6.5ft. 100 33 39 LvC3 5.24 1.2% > 6.5ft Well drained IVe slopes, severely eroded LrB2 LeRoy silt loam, 2 to 6 percent 2.66 0.6% 5.8ft. > 6.5ft Well drained lle 95 33 42 slopes, eroded JuA Juneau silt loam, 0 to 2 percent 1.46 0.3% Oft. > 6.5ft Moderately well 145 48 50 slopes drained MsC2 1.19 Ille 115 44 Mendota silt loam, 6 to 12 percent 0.3% 5.8ft. > 6.5ft Well drained 38 slopes, eroded Plano silt loam, till substratum, 0 0.62 0.1% 3.8ft. > 6.5ft PsA Well drained to 2 percent slopes



MsB	Mendota silt loam, 2 to 6 percent slopes	0.46	0.1%	5.8ft.	> 6.5ft.	Well drained	lle	125	41	48
LvD2	Lomira silt loam, 12 to 20 percent slopes, eroded	0.29	0.1%	> 6.5ft.	> 6.5ft.	Well drained	IVe	97	32.3	38
LrC2	LeRoy silt loam, 6 to 12 percent slopes, eroded	0.27	0.1%	5.8ft.	> 6.5ft.	Well drained	IIIe	85	28	38
		127	42.4	40.4						

\*c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin County: Fond Du Lac Location: 14-16N-15E Township: Rosendale Acres: 437.51



Date:



8/28/2017





Soils data provided by USDA and NRCS.

	. ,										
Area	Symbol: WI039, Soil Area V	/ersion·	16								
	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
PnA	Pella silty clay loam, cool, 0 to 2 percent slopes	184.11	42.1%		Oft.	> 6.5ft.	Poorly drained	llw	162	53.5	66.5
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	102.20	23.4%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
LvB	Lomira silt loam, 2 to 6 percent slopes	71.10	16.3%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
PhA	Pella silt loam, 0 to 2 percent slopes	32.38	7.4%		Oft.	> 6.5ft.	Poorly drained	llw	124.4	46	
VgA	Virgil silt loam, 0 to 2 percent slopes	15.55	3.6%		2ft.	> 6.5ft.	Somewhat poorly drained	llw	155	51	48
LtC3	LeRoy soils, 6 to 12 percent slopes, severely eroded	12.75	2.9%		5.8ft.	> 6.5ft.	Well drained	IVe	80	26	36
Pc	Palms mucky peat, 0 to 2 percent slopes	7.93	1.8%		Oft.	> 6.5ft.	Very poorly drained	IIIw	120		
ScA	St. Charles silt loam, 0 to 2 percent slopes	6.84	1.6%		4ft.	> 6.5ft.	Well drained	lw	150	48	51
LvA	Lomira silt loam, 0 to 2 percent slopes	2.35	0.5%		> 6.5ft.	> 6.5ft.	Well drained	lw	120	40	47
LrB2	LeRoy silt loam, 2 to 6 percent slopes, eroded	2.30	0.5%		5.8ft.	> 6.5ft.	Well drained	lle	95	33	42
	•	-	•			•	Weight	ed Average	136.4	44.7	49.8

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method

### **Tillable Soils**





State: Wisconsin County: Fond du Lac Location: 17-15N-16E Township: Lamartine

Acres: 38.56 12/11/2022 Date:









Solis	data provided by USD	A and N	RUS.						3		3	3
Are	a Symbol: WI039, S	Soil Area	a Version: 2	23			•					
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Soil Drainage	Non-Irr Class *c	Corn Bu	Soybeans Bu	Soybeans Tons	*n NCCPI Overall	*n NCCPI Corn	*n NCCPI Soybeans
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	30.37	78.8%		Well drained		115	38		75	75	67
PhA	Pella silt loam, 0 to 2 percent slopes	7.05	18.3%		Poorly drained	11\\\\	124.4		46	72	72	52
Hu	Houghton mucky peat, 0 to 2 percent slopes	0.58	1.5%		Very poorly drained	IIIw	135	46		10	10	3
BsA	Brookston silt loam, 0 to 2 percent slopes	0.49	1.3%		Poorly drained	11\4/	120	40		76	76	46
Pc	Palms mucky peat, 0 to 2 percent slopes	0.07	0.2%		Very poorly drained	IIIw	120			36	36	28
				Weighte	d Average	2.02	117.1	31.1	8.4	*n 73.4	*n 73.4	*n 63

<sup>\*</sup>n: The aggregation method is "Weighted Average using all components" \*c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin Fond Du Lac County: Location: 8-16N-14E

Township: Ripon Acres: 249.73 8/28/2017 Date:









Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
KuA	Kibbie silt loam, 0 to 2 percent slopes	41.17	16.5%		1ft.	> 6.5ft.	Somewhat poorly drained	llw	115	38	42
PhA	Pella silt loam, 0 to 2 percent slopes	40.18	16.1%		Oft.	> 6.5ft.	Poorly drained	llw	124.4	46	
PsB	Plano silt loam, till substratum, 2 to 6 percent slopes	35.37	14.2%		3.8ft.	> 6.5ft.	Well drained	lle			
MsB	Mendota silt loam, 2 to 6 percent slopes	27.96	11.2%		5.8ft.	> 6.5ft.	Well drained	lle	125	41	48
MsB2	Mendota silt loam, 2 to 6 percent slopes, eroded	19.56	7.8%		5.8ft.	> 6.5ft.	Well drained	IIIe	125	41	48
LvC2	Lomira silt loam, 6 to 12 percent slopes, eroded	17.07	6.8%		> 6.5ft.	> 6.5ft.	Well drained	IIIe	106.2	35.2	41.5
MsC2	Mendota silt loam, 6 to 12 percent slopes, eroded	15.71	6.3%		5.8ft.	> 6.5ft.	Well drained	IIIe	115	38	44
WsB	Washtenaw silt loam, 2 to 6 percent slopes	14.38	5.8%		2ft.	> 6.5ft.	Somewhat poorly drained	lle	130	43	
VgB	Virgil silt loam, 2 to 6 percent slopes	9.36	3.7%		2ft.	> 6.5ft.	Somewhat poorly drained	llw	155	51	48
WsA	Washtenaw silt loam, 0 to 2 percent slopes	6.37	2.6%		Oft.	> 6.5ft.	Poorly drained	llw	130	43	
MdC2	Markesan silt loam, 6 to 12 percent slopes, eroded	5.64	2.3%		> 6.5ft.	> 6.5ft.	Well drained	IIIe	95	32	41
PsA	Plano silt loam, till substratum, 0 to 2 percent slopes	5.04	2.0%		3.8ft.	> 6.5ft.	Well drained	I			
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	4.77	1.9%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
JuA	Juneau silt loam, 0 to 2 percent slopes	2.57	1.0%		Oft.	> 6.5ft.	Moderately well drained	I	145	48	50
BsA	Brookston silt loam, 0 to 2 percent slopes	2.20	0.9%		0.7ft.	> 6.5ft.	Poorly drained	llw	120	40	



	Markesan silt loam, 12 to 20 percent slopes, severely eroded	1.94	0.8%	5.8ft.	> 6.5ft.	Well drained	Vle	80	27	35
Рс	Palms mucky peat, 0 to 2 percent slopes	0.44	0.2%	Oft.	> 6.5ft.	Very poorly drained	IIIw	120		
						Weighte	ed Average	101.6	34.3	26

\*c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin
County: Fond Du Lac
Location: 19-16N-16E
Township: Eldorado
Acres: 249.01
Date: 8/22/2017









Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Soil Drainage	Non-Irr Class *c	Alfalfa hay	Corn	Corn silage	Oats	Soybeans	Winter wheat
PnA	Pella silty clay loam, cool, 0 to 2 percent slopes	51.38	20.6%	Ü	Oft.	Poorly drained	Ilw	liay	162	25	87	53.5	66.5
PhA	Pella silt loam, 0 to 2 percent slopes	48.20	19.4%		Oft.	Poorly drained	llw		124.4	20		46	
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	43.75	17.6%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45
LvB	Lomira silt loam, 2 to 6 percent slopes	30.60	12.3%		> 6.5ft.	Well drained	lle	4.4	115	19	75	38	45
ScA	St. Charles silt loam, 0 to 2 percent slopes	24.17	9.7%		4ft.	Well drained	lw	5.5	150	25	85	48	51
FsB	Fox silt loam, 2 to 6 percent slopes	18.04	7.2%		> 6.5ft.	Well drained	lle	4.1	105	17	67.2	35	44.8
VgA	Virgil silt loam, 0 to 2 percent slopes	10.04	4.0%		2ft.	Somewhat poorly drained	llw	5.4	155	25	80	51	48
Pc	Palms mucky peat, 0 to 2 percent slopes	8.23	3.3%		Oft.	Very poorly drained	IIIw		120	20			
ScB	St. Charles silt loam, 2 to 6 percent slopes	4.99	2.0%		4ft.	Well drained	lle	5.3	145	24	85	48	51
FsB2	Fox silt loam, 2 to 6 percent slopes, eroded	4.35	1.7%		> 6.5ft.	Well drained	lle	4.1	105	17	70	35	42
FsA	Fox silt loam, 0 to 2 percent slopes	3.29	1.3%		> 6.5ft.	Well drained	lls	4.1	111	17	67		47
LvC2	Lomira silt loam, 6 to 12 percent slopes, eroded	1.78	0.7%		> 6.5ft.	Well drained	Ille	4	106.2	17.2	66.2	35.2	41.5
VgB	Virgil silt loam, 2 to 6 percent slopes	0.19	0.1%		2ft.	Somewhat poorly drained	llw	5.4	155	25	80	51	48
		•	-		_	Weighte	d Average	2.6	131.3	21.2	61	42.4	40

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method





State: Wisconsin
County: Fond Du Lac
Location: 31-15N-15E
Township: Springvale
Acres: 157.06
Date: 8/28/2017







Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
LrB2	LeRoy silt loam, 2 to 6 percent slopes, eroded	40.82	26.0%		5.8ft.	> 6.5ft.	Well drained	lle	95	33	42
LtC3	LeRoy soils, 6 to 12 percent slopes, severely eroded	28.43	18.1%		5.8ft.	> 6.5ft.	Well drained	IVe	80	26	36
LvB2	Lomira silt loam, 2 to 6 percent slopes, eroded	28.43	18.1%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
VgA	Virgil silt loam, 0 to 2 percent slopes	12.67	8.1%		2ft.	> 6.5ft.	Somewhat poorly drained	llw	155	51	48
PhA	Pella silt loam, 0 to 2 percent slopes	11.42	7.3%		Oft.	> 6.5ft.	Poorly drained	llw	124.4	46	
LvB	Lomira silt loam, 2 to 6 percent slopes	10.95	7.0%		> 6.5ft.	> 6.5ft.	Well drained	lle	115	38	45
LvC3	Lomira silt loam, 6 to 12 percent slopes, severely eroded	10.41	6.6%		> 6.5ft.	> 6.5ft.	Well drained	IVe	100	33	39
LvA	Lomira silt loam, 0 to 2 percent slopes	7.36	4.7%		> 6.5ft.	> 6.5ft.	Well drained	lw	120	40	47
HoD3	Hochheim soils, 12 to 20 percent slopes, severely eroded	1.80	1.1%		5.8ft.	> 6.5ft.	Well drained	Vle	75	25	29
PsA	Plano silt loam, till substratum, 0 to 2 percent slopes	1.78	1.1%		3.8ft.	> 6.5ft.	Well drained	I			
LtD3	LeRoy soils, 12 to 20 percent slopes, severely eroded	1.32	0.8%		5.8ft.	> 6.5ft.	Well drained	Vle	70	23	32
BsA	Brookston silt loam, 0 to 2 percent slopes	1.11	0.7%		0.7ft.	> 6.5ft.	Poorly drained	llw	120	40	
HmD	Hochheim loam, 12 to 20 percent slopes	0.56	0.4%		> 6.5ft.	1.2ft. (Densic material)	Well drained	IVe			
	Weighted Average 10										

<sup>\*</sup>c: Using Capabilities Class Dominant Condition Aggregation Method