



Soils Map



Soils data provided by USDA and NRCS.

©2017 AgriData, Inc.



State: **Indiana**
 County: **Warren**
 Location: **12-22N-8W**
 Township: **Liberty**
 Acres: **224.59**
 Date: **2/10/2017**



© AgriData, Inc. 2017 www.AgriDataInc.com



Area Symbol: IN171, Soil Area Version: 19

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat
MpC3	Miami clay loam, 6 to 12 percent slopes, severely eroded	61.11	27.2%		2.5ft.	2.8ft. (Densic material)	Moderately well drained	IVe	121	41	54
RfB2	Rainsville-Williamstown-Rockfield silt loams, 2 to 6 percent slopes, eroded	59.13	26.3%		2.2ft.	4.6ft. (Densic material)	Moderately well drained	Ile	133	47	60
Cz	Cyclone silty clay loam, 0 to 2 percent slopes	33.53	14.9%		0.2ft.	> 6.5ft.	Poorly drained	IIlw	184	54	75
RoB2	Rockfield silt loam, 2 to 6 percent slopes, eroded	24.58	10.9%		2.2ft.	4.6ft. (Densic material)	Moderately well drained	Ile	141	50	64
SIA	Starks silt loam, till substratum, 0 to 2 percent slopes	20.82	9.3%		1.2ft.	4.6ft. (Densic material)	Somewhat poorly drained	IIlw	160	52	72
Wh	Washtenaw silt loam	10.02	4.5%		0.2ft.	> 6.5ft.	Poorly drained	IIlw	165	49	66
PuB2	Proctor silt loam, till substratum, 2 to 6 percent slopes, eroded	5.07	2.3%		2.2ft.	> 6.5ft.	Moderately well drained	Ile	150	50	68
StB3	Strawn clay loam, 2 to 6 percent slopes, severely eroded	5.05	2.2%		1.5ft.	1.7ft. (Densic material)	Moderately well drained	IIIe	131	45	59
Dw	Drummer silty clay loams	4.74	2.1%		0.5ft.	> 6.5ft.	Poorly drained	IIlw	175	49	70
Bk	Beckville loam, occasionally flooded	0.54	0.2%		2.2ft.	> 6.5ft.	Moderately well drained	IIlw			
Weighted Average									143.1	47.2	62.6

Area Symbol: IN171, Soil Area Version: 19

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.