



# Soils Map



State: **Ohio**  
 County: **Union**  
 Location: **40° 28' 1.9, -83° 21' 24.86**  
 Township: **Jackson**  
 Acres: **621.81**  
 Date: **3/2/2023**



Maps Provided By:



Soils data provided by USDA and NRCS.

©2023 AgriData, Inc.

© AgriData, Inc. 2023

www.AgriDataInc.com

Area Symbol: OH159, Soil Area Version: 21

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn Bu	Soybeans Bu	Wheat Bu	Winter wheat Bu	*n NCCPI Corn	*n NCCPI Soybeans	*eFOTG R
Blg1A1	Blount silt loam, ground moraine, 0 to 2 percent slopes	300.49	48.3%		0.7ft.	3.2ft. (Densic material)	Somewhat poorly drained	llw	141	45		63	70	58	78
We	Wetzel silty clay loam	111.66	18.0%		0.8ft.	> 6.5ft.	Poorly drained	llw	145	51		47.3	84	72	78
Blg1B1	Blount silt loam, ground moraine, 2 to 4 percent slopes	100.18	16.1%		0.7ft.	3.1ft. (Densic material)	Somewhat poorly drained	lle	137	44		61	68	55	75
Mns3A	Minster silty clay loam, 0 to 1 percent slopes	56.78	9.1%		0.5ft.	> 6.5ft.	Very poorly drained	lllw	158	47		61	52	44	75
FoA	Fox silt loam, till plain, 0 to 2 percent slopes	18.15	2.9%		> 6.5ft.	2.7ft. (Strongly contrasting textural stratification)	Well drained	lls	102	37		51	65	59	71
Ee	Eel silt loam, 0 to 2 percent slopes, occasionally flooded	8.86	1.4%		1.6ft.	> 6.5ft.	Moderately well drained	llw	123	43	8		81	92	71

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn Bu	Soybeans Bu	Wheat Bu	Winter wheat Bu	*n NCCPI Corn	*n NCCPI Soybeans	*eFOTG PI
So	Sloan silty clay loam, 0 to 1 percent slopes, frequently flooded	7.65	1.2%		0.3ft.	> 6.5ft.	Very poorly drained	Illw	134	47			74	70	78
Lc	Lippincott silty clay loam, 0 to 2 percent slopes	7.35	1.2%		0.3ft.	3ft. (Strongly contrasting textural stratification)	Very poorly drained	llw	127	44		57	58	57	80
Gwg1B1	Glynwood silt loam, ground moraine, 2 to 6 percent slopes	7.03	1.1%		1.5ft.	2.8ft. (Densic material)	Moderately well drained	lle	132	46		60	67	53	70
Gwg5C2	Glynwood clay loam, ground moraine, 6 to 12 percent slopes, eroded	1.85	0.3%		1.5ft.	2.4ft. (Densic material)	Moderately well drained	lle	117	29		53	59	36	61
HeA	Henshaw silt loam, 0 to 2 percent slopes	1.17	0.2%		1ft.	> 6.5ft.	Somewhat poorly drained	llw	140	45		66	82	68	73
FoC2	Fox silt loam, 6 to 12 percent slopes, moderately eroded	0.64	0.1%		> 6.5ft.	2.6ft. (Strongly contrasting textural stratification)	Well drained	lle	98			47	64	59	59
<b>Weighted Average</b>								<b>2.11</b>	<b>140.8</b>	<b>45.8</b>	<b>0.1</b>	<b>57.5</b>	<b>*n 70.4</b>	<b>*n 59.3</b>	<b>76.8</b>

\*eftog PI: Obtained from the NRCS eFOTG (<http://efotg.sc.egov.usda.gov>)

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method