
Soils Map





State: **Georgia** County: **Sumter**

Location: 32° 0' 15.54, -84° 0' 6.02

Township: Leslie-De Soto

Acres: **6916.85**Date: **4/14/2023**







Soils data provided by USDA and NRCS.

eroded

Collis data provided by Cobh and Nicoo.															
Area	Area Symbol: GA652, Soil Area Version: 18														
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non- Irr Class *c	Irr Class *c	Corn Bu	Corn Irrigated Bu	Cotton lint Lbs	Cotton lint Irrigated Lbs	COybcans	Soybeans Irrigated Bu
GoB	Greenville sandy loam, 2 to 5 percent slopes	2420.26	35.0%		> 6.5ft.	> 6.5ft.	Well drained	lle							
Grd	Grady soils	831.34	12.0%		Oft.	> 6.5ft.	Poorly drained	Vw							
lg	Irvington sandy loam	546.36	7.9%		1.5ft.	> 6.5ft.	Somewhat poorly drained	llw							
Kib	Kinston and Bibb soils, 0 to 2 percent slopes, frequently flooded	330.13	4.8%		0.5ft.	> 6.5ft.	Poorly drained	VIw							
LMB	Lucy loamy sand, 0 to 5 percent slopes	320.01	4.6%		> 6.5ft.	> 6.5ft.	Well drained	lls	lls	75		550			
GqC3	Greenville sandy clay loam, 5 to 8 percent slopes, severely	282.33	4.1%		> 6.5ft.	> 6.5ft.	Well drained	lVe							

© AgriData	, Inc. 2023 v	www.AgriDataIn	c.com	•			1		•						
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non- Irr Class *c	Irr Class *c	Corn Bu	Corn Irrigated Bu	Cotton lint Lbs	Cotton lint Irrigated Lbs	Soybeans Bu	Soybeans Irrigated Bu
GoA	Greenville sandy loam, 0 to 2 percent slopes	253.76	3.7%		> 6.5ft.	> 6.5ft.	Well drained	1							
OeB	Orangeburg loamy sand, 2 to 5 percent slopes	242.63	3.5%		> 6.5ft.	> 6.5ft.	Well drained	lle	lle	113.3		966.7		41.7	
Gt	Goldsboro loamy sand, 0 to 2 percent slopes	240.56	3.5%		2.3ft.	> 6.5ft.	Moderately well drained	llw		125		700		42	
TuB2	Tifton sandy loam, 2 to 5 percent slopes, moderately eroded	217.04	3.1%		3.5ft.	> 6.5ft.	Well drained	lle							
RhB	Red Bay sandy loam, 2 to 5 percent slopes	207.31	3.0%		> 6.5ft.	> 6.5ft.	Well drained	lle							
TuA	Tifton sandy loam, 0 to 2 percent slopes	187.92	2.7%		3.5ft.	> 6.5ft.	Well drained	I							
GoC2	Greenville sandy loam, 5 to 8 percent slopes, eroded	140.71	2.0%		> 6.5ft.	> 6.5ft.	Well drained	llle							
Oi	Ochlockonee soils, local alluvium	114.07	1.6%		4ft.	> 6.5ft.	Well drained	llw							
	Faceville sandy loam, 2 to 5 percent slopes, moderately eroded	113.20	1.6%		> 6.5ft.	> 6.5ft.	Well drained	llle							
CoC2	Carnegie sandy loam, 5 to 8 percent slopes, moderately eroded	55.77	0.8%		> 6.5ft.	> 6.5ft.	Well drained	IVe	llle						
CoB2	Carnegie sandy loam, 2 to 5 percent slopes, moderately eroded	53.72	0.8%		> 6.5ft.	> 6.5ft.	Well drained	∭e							

e / igiibata	, Inc. 2023 \	www.AgriDatain	0.00111				1								
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non- Irr Class *c	Irr Class *c	Corn Bu	Corn Irrigated Bu	Cotton lint Lbs	Cotton lint Irrigated Lbs	Soybeans Bu	Soybeans Irrigated Bu
CoD2	Carnegie sandy loam, 8 to 12 percent slopes, moderately eroded	46.08	0.7%		> 6.5ft.	> 6.5ft.	Well drained	Vle	IVe						
HdC	Henderson cherty sandy loam, 2 to 8 percent slopes	40.13	0.6%		> 6.5ft.	> 6.5ft.	Well drained	lVe							
NhB	Norfolk loamy sand, 2 to 5 percent slopes	38.75	0.6%		5ft.	> 6.5ft.	Well drained	lle	e	99.8		703.2		35.6	
RhC2	Red Bay sandy loam, 5 to 8 percent slopes, eroded	34.54	0.5%		> 6.5ft.	> 6.5ft.	Well drained	lVe							
GqD3	Greenville sandy clay loam, 8 to 12 percent slopes, severely eroded	34.09	0.5%		> 6.5ft.	> 6.5ft.	Well drained	Vle							
LMC	Lucy loamy sand, 5 to 8 percent slopes	29.88	0.4%		> 6.5ft.	> 6.5ft.	Well drained	IIIs	<u>Is</u>	70		600		25	
NhA	Norfolk loamy sand, 0 to 2 percent slopes	29.42	0.4%		5ft.	> 6.5ft.	Well drained	I	I	110	190	735.7	1000	39.3	55
W	Water	18.22	0.3%		> 6.5ft.	> 6.5ft.									
RhA	Red Bay sandy loam, 0 to 2 percent slopes	15.23	0.2%		> 6.5ft.	> 6.5ft.	Well drained	I							
HdE	Henderson cherty sandy loam, 8 to 17 percent slopes	12.10	0.2%		> 6.5ft.	> 6.5ft.	Well drained	VIIe							
TuC2	Tifton sandy loam, 5 to 8 percent slopes, eroded	11.06	0.2%		3.5ft.	> 6.5ft.	Well drained	llle							
OeA	Orangeburg loamy sand, 0 to 2 percent slopes	10.80	0.2%		> 6.5ft.	> 6.5ft.	Well drained	ı	ı	117.6	190	900	1100	45	55



© AgriData	⊌ AgriData, Inc. 2023 www.AgriDataInc.com														
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non- Irr Class *c	Irr Class *c	Corn Bu	Corn Irrigated Bu	Cotton lint Lbs	Cotton lint Irrigated Lbs	Soybeans Bu	Soybeans Irrigated Bu
OeC2	Orangeburg loamy sand, 5 to 8 percent slopes, moderately eroded	10.68	0.2%		> 6.5ft.	> 6.5ft.	Well drained	≡ e		113.3		966.7		41.7	
OeD2	Orangeburg loamy sand, 8 to 12 percent slopes, moderately eroded	10.25	0.1%		> 6.5ft.	> 6.5ft.	Well drained	IVe							
FuC2	Faceville sandy loam, 5 to 8 percent slopes, moderately eroded	5.70	0.1%		> 6.5ft.	> 6.5ft.	Well drained	≡ e							
ArB	Americus loamy sand, 0 to 5 percent slopes	4.74	0.1%		> 6.5ft.	> 6.5ft.	Somewhat excessively drained	<u> </u>							
LMD	Lucy loamy sand, 8 to 15 percent slopes	4.47	0.1%		> 6.5ft.	> 6.5ft.	Well drained	IVs							
Ros	Rains sandy loam	2.54	0.0%		0.5ft.	> 6.5ft.	Poorly drained	VIw							
FuA	Faceville sandy loam, 0 to 2 percent slopes	1.05	0.0%		> 6.5ft.	> 6.5ft.	Well drained	I							
	Weighted Average									13.5	1.1	96.3	6	3.5	0.3
<u> </u>															

^{*}n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

*- Non Irr Class weighted average cannot be calculated on the current soils data due to missing data.

*- Irr Class weighted average cannot be calculated on the current soils data due to missing data.