



Soils Map



Soils data provided by USDA and NRCS.

©2017 AgriData, Inc.



State: **Ohio**
 County: **Pike**
 Location: **39° 9' 25.4, -82° 53' 13.7**
 Township: **Jackson**
 Acres: **1769.02**
 Date: **9/18/2017**



© AgriData, Inc. 2017 www.AgriDataInc.com



Area Symbol: OH131, Soil Area Version: 19
 Area Symbol: OH141, Soil Area Version: 17

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat	*eFOTG PI
Mh	Martinsville loam, rarely flooded	474.48	26.8%		> 6.5ft.	> 6.5ft.	Well drained	I	130	45	50	81
En	Elkinsville silt loam, rarely flooded	428.46	24.2%		> 6.5ft.	> 6.5ft.	Well drained	I	130	42	48	78
FoA	Fox loam, 0 to 2 percent slopes	170.89	9.7%		> 6.5ft.	> 6.5ft.	Well drained	IIs	110	32	45	71
FoB	Fox loam, 2 to 6 percent slopes	163.93	9.3%		> 6.5ft.	> 6.5ft.	Well drained	Ile	95	30	42	68
Ge	Genesee silt loam, occasionally flooded	125.05	7.1%		> 6.5ft.	> 6.5ft.	Well drained	IIw	130	37	35	71
Th	Taggart silt loam, rarely flooded	106.79	6.0%		2ft.	> 6.5ft.	Somewhat poorly drained	IIw	115	40	40	70
FoC	Fox loam, 6 to 12 percent slopes	104.22	5.9%		> 6.5ft.	> 6.5ft.	Well drained	IIle	85	28	38	61
PaA	Parke silt loam, 0 to 3 percent slopes	71.44	4.0%		> 6.5ft.	> 6.5ft.	Well drained	I	125	45	50	76
St	Stonelick loam, occasionally flooded	36.85	2.1%		> 6.5ft.	> 6.5ft.	Well drained	IIw	95	30	30	62
PaB	Parke silt loam, 3 to 8 percent slopes	27.13	1.5%		> 6.5ft.	> 6.5ft.	Well drained	Ile	115	40	46	74
Or	Orrville silt loam, 0 to 3 percent slopes, frequently flooded	15.08	0.9%		1ft.	> 6.5ft.	Somewhat poorly drained	IIw	109	35	48	69
TbA	Taggart silt loam, 0 to 2 percent slopes	14.27	0.8%		1.2ft.	> 6.5ft.	Somewhat poorly drained	IIw	130	45	45	70
LrB	Libre silt loam, 2 to 6 percent slopes	7.08	0.4%		2.7ft.	> 6.5ft.	Moderately well drained	Ile	135	45	45	74



NgC	Negley loam, 8 to 15 percent slopes	6.91	0.4%		> 6.5ft.	> 6.5ft.	Well drained	IIIe	90	30	40	59
NgE	Negley loam, 25 to 35 percent slopes	5.94	0.3%		> 6.5ft.	> 6.5ft.	Well drained	VIe				0
RbA	Rainsboro silt loam, 0 to 2 percent slopes	4.96	0.3%		2.7ft.	2.2ft. (Fragipan)	Moderately well drained	IIw	130	40	45	71
MaD2	Markland silty clay loam, 15 to 25 percent slopes, eroded	1.62	0.1%		3.9ft.	> 6.5ft.	Well drained	VIe				50
Pg	Pits, gravel	1.55	0.1%		> 6.5ft.	> 6.5ft.						0
FoC2	Fox loam, 6 to 12 percent slopes, eroded	1.15	0.1%		> 6.5ft.	2.5ft. (Strongly contrasting textural stratification)	Well drained	IIIe	110	30	40	59
NeE2	Negley loam, 20 to 35 percent slopes, eroded	0.58	0.0%		> 6.5ft.	> 6.5ft.	Well drained	VIe				0
MaB2	Markland silty clay loam, 3 to 8 percent slopes, eroded	0.37	0.0%		3.9ft.	> 6.5ft.	Well drained	IIIe	100	28	36	62
MkA	McGary silt loam, 0 to 4 percent slopes	0.27	0.0%		2ft.	> 6.5ft.	Somewhat poorly drained	IIw	100	30	35	67
Weighted Average									119.1	38.9	45	74.1

*eftog PI: Obtained from the NRCS eFOTG (<http://efotg.sc.egov.usda.gov>)

*eftog PI index for OH was updated on 3/8/2017

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.