



Soils Map



State: **Indiana**
 County: **Jasper**
 Location: **10-28N-6W**
 Township: **Milroy**
 Acres: **417.01**
 Date: **6/26/2017**



Soils data provided by USDA and NRCS.

Area Symbol: IN073, Soil Area Version: 17

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Restrictive Layer | Soil Drainage | Non-Irr Class *c | Corn | Corn Irrigated | Soybeans | Soybeans Irrigated | Winter wheat |
|------|---|-------|------------------|----------------------|-------------|--------------------------|-------------------------|------------------|------|----------------|----------|--------------------|--------------|
| Gf | Gilford fine sandy loam | 72.16 | 17.3% | | 0.2ft. | > 6.5ft. | Poorly drained | Ilw | 148 | 6 | 33 | | 59 |
| OaB | Oakville fine sand, 2 to 6 percent slopes | 60.18 | 14.4% | | > 6.5ft. | > 6.5ft. | Well drained | IVs | 73 | | 25 | | 36 |
| Wm | Watseka-Maumee loamy sands | 51.28 | 12.3% | | 0.5ft. | > 6.5ft. | Poorly drained | IIIs | 125 | | 29 | | 53 |
| Mu | Morocco loamy sand | 48.60 | 11.7% | | 1.2ft. | > 6.5ft. | Somewhat poorly drained | IIIs | 100 | | 29 | | 45 |
| Za | Zadog-Maumee loamy sands | 47.15 | 11.3% | | 0.2ft. | > 6.5ft. | Poorly drained | IIIw | 146 | | 36 | | 58 |
| BeB | Brems loamy sand, 1 to 3 percent slopes | 31.58 | 7.6% | | 2.5ft. | > 6.5ft. | Moderately well drained | IVs | 91 | 5 | 32 | 2 | 41 |
| Ir | Iroquois fine sandy loam | 21.97 | 5.3% | | 0.2ft. | > 6.5ft. | Very poorly drained | Ilw | 170 | | 47 | | 68 |
| Pa | Papineau sandy loam | 21.30 | 5.1% | | 1.2ft. | > 6.5ft. | Somewhat poorly drained | Ilw | 150 | | 46 | | 68 |
| Px | Prochaska loamy sand, frequently flooded | 18.05 | 4.3% | | 0.2ft. | > 6.5ft. | Very poorly drained | IIIw | 110 | | 19 | | |
| SmA | Simonin loamy sand, 0 to 2 percent slopes | 17.59 | 4.2% | | 2.2ft. | 3.3ft. (Densic material) | Moderately well drained | IIs | 131 | | 43 | | 59 |
| Re | Rensselaer loam | 14.53 | 3.5% | | 0.7ft. | > 6.5ft. | Poorly drained | Ilw | 172 | | 48 | | 69 |
| Fa | Faxon loam | 5.36 | 1.3% | | 0.2ft. | 3ft. (Lithic bedrock) | Very poorly drained | Ilw | 145 | | 39 | | 58 |
| We | Watseka loamy fine sand | 4.32 | 1.0% | | 1.2ft. | > 6.5ft. | Somewhat poorly drained | IIIs | 105 | | 27 | | 47 |



| | | | | | | | | | | | | | |
|-------------------------|--|------|------|--|----------|----------|-------------------------|-----|--------------|------------|-------------|------------|-------------|
| NaB | Nesius fine sand, 1 to 3 percent slopes | 2.78 | 0.7% | | 2.2ft. | > 6.5ft. | Moderately well drained | IVs | 95 | | 30 | | 43 |
| OaC | Oakville fine sand, 6 to 15 percent slopes | 0.16 | 0.0% | | > 6.5ft. | > 6.5ft. | Well drained | Vle | 64 | | 23 | | 32 |
| Weighted Average | | | | | | | | | 123.1 | 1.4 | 32.9 | 0.2 | 50.3 |

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.