



Soils Map



State: **Michigan**
 County: **Calhoun**
 Location: **8-4S-5W**
 Township: **Clarendon**
 Acres: **161.09**
 Date: **6/8/2017**



Maps Provided By:



© AgriData, Inc. 2017

www.AgriDataInc.com



Soils data provided by USDA and NRCS.

Archived Soils Ending 11/6/2014

Area Symbol: MI025, Soil Area Version: 10

| Cod e | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Soil Drainage | Non-Irr Class *c | Irr Class *c | Alfalfa hay | Corn | Corn Irrigated | Corn silage | Corn silage Irrigated | Oats | Soybeans | Soybeans Irrigated | Winter wheat | Winter wheat Irrigated |
|-------|--|-------|------------------|----------------------|-------------|-------------------------|------------------|--------------|-------------|------|----------------|-------------|-----------------------|------|----------|--------------------|--------------|------------------------|
| 65 | Sebewa loam | 38.16 | 23.7% | | 0ft. | Poorly drained | Ilw | | 5 | 108 | | 20 | | 94 | 38 | | 54 | |
| 33B | Riddles loam, 0 to 6 percent slopes | 32.79 | 20.4% | | > 6.5ft. | Well drained | Ile | | 5 | 112 | 166 | 20 | 26 | 8 | 37 | 62 | 58 | |
| 25B | Kalamazoo loam, 2 to 6 percent slopes | 31.00 | 19.2% | | > 6.5ft. | Well drained | Ile | Ile | 4 | 99 | 160 | 17 | 25 | 76 | 30 | 55 | 39 | |
| 21B | Leoni gravelly loam, 0 to 6 percent slopes | 23.49 | 14.6% | | > 6.5ft. | Well drained | Ills | | 5 | 109 | 160 | 18 | 27 | 80 | 34 | 54 | 54 | 36 |
| 45A | Sleeth loam, 0 to 2 percent slopes | 18.36 | 11.4% | | 2ft. | Somewhat poorly drained | Ilw | | 5 | 121 | | 21 | | 8 | 42 | | 64 | |
| 33C | Riddles loam, 6 to 12 percent slopes | 9.23 | 5.7% | | > 6.5ft. | Well drained | Ille | | 5 | 106 | 152 | 18 | 25 | 8 | 33 | 57 | 52 | |
| 25C | Kalamazoo loam, 6 to 12 percent slopes | 5.07 | 3.1% | | > 6.5ft. | Well drained | Ille | | 4 | 89 | 145 | 16 | 24 | 71 | 27 | 50 | 36 | |
| 43B | Brady sandy loam, 1 to 4 percent slopes | 1.73 | 1.1% | | 2ft. | Somewhat poorly drained | Ile | | 5 | 106 | | 20 | | 85 | 33 | | 49 | |





| | | | | | | | | | | | | | | | | | | |
|-------------------------|--|------|------|--|----------|--------------|------|------|------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|------------|
| 13B | Spinks loamy sand, 0 to 6 percent slopes | 1.26 | 0.8% | | > 6.5ft. | Well drained | Ills | Ills | 4 | 77 | 165 | 13 | 24 | 7 | 27 | 50 | 31 | |
| Weighted Average | | | | | | | | | 4.8 | 107.7 | 102.5 | 18.9 | 16.4 | 54.8 | 35.4 | 36.3 | 52.2 | 5.2 |

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.

Soils Map



State: **Michigan**
 County: **Calhoun**
 Location: **20-4S-6W**
 Township: **Tekonsha**
 Acres: **217.83**
 Date: **6/8/2017**



Soils data provided by USDA and NRCS.

Archived Soils Ending 11/6/2014

Area Symbol: MI025, Soil Area Version: 10

| Cod e | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Soil Drainage | Non-Irr Class *c | Irr Class *c | Alfalfa hay | Corn | Corn Irrigated | Corn silage | Corn silage Irrigated | Oats | Soybeans | Soybeans Irrigated | Winter wheat |
|-------------------------|---|-------|------------------|----------------------|-------------|-------------------------|------------------|--------------|-------------|--------------|----------------|-------------|-----------------------|-------------|-------------|--------------------|--------------|
| 43B | Brady sandy loam, 1 to 4 percent slopes | 72.76 | 33.4% | | 2ft. | Somewhat poorly drained | Ile | | 5 | 106 | | 20 | | 85 | 33 | | 49 |
| 14B | Bronson sandy loam, 0 to 6 percent slopes | 53.37 | 24.5% | | 2.7ft. | Moderately well drained | Ile | Ils | 4 | 95 | 155 | 16 | 26 | 74 | 27 | 53 | 35 |
| 16B | Oshtemo sandy loam, 0 to 6 percent slopes | 50.43 | 23.2% | | > 6.5ft. | Well drained | IIIs | IIle | 4 | 100 | 165 | 16 | 27 | 80 | 30 | 55 | 45 |
| 44A | Matherton loam, 0 to 3 percent slopes | 20.66 | 9.5% | | 1.5ft. | Somewhat poorly drained | IIw | | 5 | 107 | | 17 | | 86 | 32 | | 40 |
| 65 | Sebewa loam | 17.52 | 8.0% | | 0ft. | Poorly drained | IIw | | 5 | 108 | | 20 | | 94 | 38 | | 54 |
| 13B | Spinks loamy sand, 0 to 6 percent slopes | 3.09 | 1.4% | | > 6.5ft. | Well drained | IIIs | IIIs | 4 | 77 | 165 | 13 | 24 | 7 | 27 | 50 | 31 |
| Weighted Average | | | | | | | | | 4.5 | 101.8 | 78.5 | 17.7 | 13 | 80.9 | 31.1 | 26.4 | 43.9 |

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.