



Soils Map



State: **Indiana**
 County: **Pulaski**
 Location: **8-31N-3W**
 Township: **Rich Grove**
 Acres: **1439.51**
 Date: **9/7/2022**



Maps Provided By:



© AgriData, Inc. 2021

www.AgrIDataInc.com



Soils data provided by USDA and NRCS.

©2022 AgriData, Inc.

Area Symbol: IN131, Soil Area Version: 23

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Restrictive Layer | Soil Drainage | Non-Irr Class *c | Corn Bu | Corn Irrigated Bu | Soybeans Bu | Soybeans Irrigated Bu | Winter wheat Bu | *n NCCPI Soybeans |
|-------|-----------------------------------------------------|--------|------------------|----------------------|-------------|-------------------|---------------------|------------------|---------|-------------------|-------------|-----------------------|-----------------|-------------------|
| MhaA | Maumee loamy fine sand, 0 to 1 percent slopes | 398.71 | 27.7% | | 0.2ft. | > 6.5ft. | Poorly drained | Illw | 134 | | 47 | | 54 | 38 |
| TmaAN | Toto muck, drained, 0 to 1 percent slopes | 281.36 | 19.5% | | 0ft. | > 6.5ft. | Very poorly drained | IVw | 136 | | 35 | | 55 | 78 |
| MvhAN | Moston muck, drained, 0 to 1 percent slopes | 209.08 | 14.5% | | 0ft. | > 6.5ft. | Very poorly drained | IVw | 136 | | 35 | | 54 | 90 |
| MhbA | Maumee mucky loamy fine sand, 0 to 1 percent slopes | 125.29 | 8.7% | | 0.2ft. | > 6.5ft. | Very poorly drained | Illw | 136 | | 28 | | 54 | 42 |
| AbhAN | Adrian muck, drained, 0 to 1 percent slopes | 86.13 | 6.0% | | 0ft. | > 6.5ft. | Very poorly drained | Illw | 144 | | 40 | | 58 | 86 |
| ReyA | Rensselaer loam, 0 to 1 percent slopes | 73.14 | 5.1% | | 0.7ft. | > 6.5ft. | Poorly drained | Ilw | 172 | | 48 | | 69 | 77 |

Soils data provided by USDA and NRCS.

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Restrictive Layer | Soil Drainage | Non-Irr Class *c | Corn Bu | Corn Irrigated Bu | Soybeans Bu | Soybeans Irrigated Bu | Winter wheat Bu | *n NCCPI Soybeans |
|-------|--------------------------------------------------------------|-------|------------------|----------------------|-------------|-------------------------------------------------------|-------------------------|------------------|---------|-------------------|-------------|-----------------------|-----------------|-------------------|
| MupA | Morocco loamy fine sand, 0 to 2 percent slopes | 72.33 | 5.0% | | 1ft. | > 6.5ft. | Somewhat poorly drained | Ills | 102 | | 36 | | 41 | 36 |
| MwzAN | Muskego muck, drained, 0 to 1 percent slopes | 53.77 | 3.7% | | 0ft. | > 6.5ft. | Very poorly drained | IVw | 138 | | 36 | | 55 | 89 |
| MgyA | Maumee-Gilford complex, 0 to 1 percent slopes | 36.62 | 2.5% | | 0.2ft. | > 6.5ft. | Poorly drained | Illw | 140 | | 30 | | 56 | 48 |
| OacB | Oakville-Denham fine sands, 1 to 5 percent slopes | 27.22 | 1.9% | | 3ft. | > 6.5ft. | Excessively drained | IVs | 84 | | 30 | | 38 | 30 |
| BstB | Brems loamy fine sand, 1 to 4 percent slopes | 20.00 | 1.4% | | 2.5ft. | > 6.5ft. | Moderately well drained | IVs | 97 | | 33 | | 44 | 34 |
| WoxB | Williamstown-Winamac fine sandy loams, 1 to 5 percent slopes | 16.45 | 1.1% | | 1.5ft. | 3.2ft. (Densic material) | Moderately well drained | Ile | 140 | | 47 | | 63 | 56 |
| EcrAN | Edselton muck, drained, 0 to 1 percent slopes | 12.14 | 0.8% | | 0ft. | > 6.5ft. | Very poorly drained | IVw | 136 | | 35 | | 54 | 84 |
| BrvA | Brady fine sandy loam, 0 to 1 percent slopes | 11.02 | 0.8% | | 1.2ft. | 4.6ft. (Strongly contrasting textural stratification) | Somewhat poorly drained | Ilw | 125 | 9 | 35 | 3 | 56 | 54 |
| BswA | Brems-Morocco loamy fine sands, 0 to 1 percent slopes | 6.72 | 0.5% | | 2.5ft. | > 6.5ft. | Moderately well drained | IVs | 98 | | 31 | | 44 | 35 |
| AatAN | Ackerman muck, drained, 0 to 1 percent slopes | 3.00 | 0.2% | | 0.2ft. | > 6.5ft. | Very poorly drained | IVw | 144 | | 35 | | 58 | 66 |
| OaeC | Oakville fine sand, 5 to 12 percent slopes | 2.75 | 0.2% | | > 6.5ft. | > 6.5ft. | Excessively drained | VIIs | 73 | | 26 | | 33 | 30 |
| ApuAN | Antung muck, drained, 0 to 1 percent slopes | 2.52 | 0.2% | | 0.2ft. | > 6.5ft. | Very poorly drained | Illw | 144 | | 35 | | 58 | 62 |





| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Water Table | Restrictive Layer | Soil Drainage | Non-Irr Class *c | Corn Bu | Corn Irrigated Bu | Soybeans Bu | Soybeans Irrigated Bu | Winter wheat Bu | *n NCCPI Soybeans |
|-------------------------|--------------------------------------------------------|-------|------------------|----------------------|-------------|-------------------|----------------|------------------|--------------|-------------------|-------------|-----------------------|-----------------|-------------------|
| NofA | Newton-Morocco loamy fine sands, 0 to 1 percent slopes | 1.26 | 0.1% | | 0.2ft. | > 6.5ft. | Poorly drained | IVw | 115 | | 28 | | 48 | 37 |
| Weighted Average | | | | | | | | 3.36 | 134.4 | 0.1 | 38.6 | *- | 54.2 | *n 61.2 |

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils Map



State: **Indiana**
 County: **Pulaski**
 Location: **18-31N-3W**
 Township: **Rich Grove**
 Acres: **179.69**
 Date: **8/7/2023**



Maps Provided By:



Soils data provided by USDA and NRCS.

©2023 AgriData, Inc.

© AgriData, Inc. 2023

www.AgriDataInc.com

Area Symbol: IN131, Soil Area Version: 24

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Non-Irr Class *c | Corn Bu | Soybeans Bu | *n NCCPI Corn | *n NCCPI Soybeans |
|-------------------------|------------------------------------------------------------------|-------|------------------|----------------------|------------------|--------------|-------------|----------------|-------------------|
| MhaA | Maumee loamy fine sand, 0 to 1 percent slopes | 64.61 | 36.0% | | IIIw | 134 | 47 | 55 | 37 |
| TmaAN | Toto muck, drained, 0 to 1 percent slopes | 39.23 | 21.8% | | IVw | 136 | 35 | 75 | 78 |
| MgyA | Maumee-Gilford complex, 0 to 1 percent slopes | 19.44 | 10.8% | | IIIw | 140 | 30 | 50 | 46 |
| GmnA | Goodell-Gilford fine sandy loams, 0 to 1 percent slopes | 16.74 | 9.3% | | IIw | 150 | 33 | 70 | 62 |
| MhbA | Maumee mucky loamy fine sand, 0 to 1 percent slopes | 13.06 | 7.3% | | IIIw | 136 | 28 | 39 | 40 |
| AbhAP | Adrian muck, drained, prairie peninsula, 0 to 1 percent slopes | 7.17 | 4.0% | | IIIw | 144 | 40 | 80 | 85 |
| MupA | Morocco loamy fine sand, 0 to 2 percent slopes | 5.52 | 3.1% | | IIIs | 102 | 36 | 51 | 34 |
| OaeC | Oakville fine sand, 5 to 12 percent slopes | 4.58 | 2.5% | | VI s | 73 | 26 | 26 | 25 |
| BstB | Brems loamy fine sand, 1 to 4 percent slopes | 3.44 | 1.9% | | IVs | 97 | 33 | 33 | 31 |
| NofA | Newton-Morocco loamy fine sands, 0 to 1 percent slopes | 2.55 | 1.4% | | IVw | 115 | 28 | 27 | 33 |
| HplrA | Houghton muck, drained, prairie peninsula, 0 to 1 percent slopes | 2.04 | 1.1% | | IIIw | 154 | 54 | 91 | 88 |
| BswA | Brems-Morocco loamy fine sands, 0 to 1 percent slopes | 1.31 | 0.7% | | IVs | 98 | 31 | 32 | 31 |
| Weighted Average | | | | | 3.24 | 133.6 | 38.1 | *n 58.6 | *n 51.4 |

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.