



Soils Map



State: **Ohio**
 County: **Lorain**
 Location: **41° 20' 41.16, -82° 11' 44.87**
 Township: **New Russia**
 Acres: **580.32**
 Date: **12/12/2017**



Soils data provided by USDA and NRCS.

©2017 AgriData, Inc.

Area Symbol: OH093, Soil Area Version: 16

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Water Table	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Corn	Soybeans	Winter wheat	*eFOTG PI
MgA	Mahoning silt loam, 0 to 2 percent slopes	301.55	52.0%		0.6ft.	> 6.5ft.	Somewhat poorly drained	IIIw				63
TrA	Trumbull silty clay loam, 0 to 2 percent slopes	110.83	19.1%		0ft.	> 6.5ft.	Poorly drained	IVw				55
Ly	Luray silty clay loam, 0 to 2 percent slopes	50.79	8.8%		0ft.	> 6.5ft.	Very poorly drained	IIw				78
FcA	Fitchville silt loam, 0 to 2 percent slopes	45.80	7.9%		0.7ft.	> 6.5ft.	Somewhat poorly drained	IIw				70
EIB2	Ellsworth silt loam, 2 to 6 percent slopes, eroded	15.29	2.6%		1ft.	> 6.5ft.	Moderately well drained	IIIe				63
Sb	Sebring silt loam, 0 to 2 percent slopes	13.78	2.4%		0.1ft.	> 6.5ft.	Poorly drained	IIIw				58
EIB	Ellsworth silt loam, 2 to 6 percent slopes	13.20	2.3%		1ft.	> 6.5ft.	Moderately well drained	IIIe				67
Or	Orrville silt loam	12.83	2.2%		1ft.	> 6.5ft.	Somewhat poorly drained	IIw	120	36	54	72
MkA	Mahoning-Tiro silt loams, 0 to 2 percent slopes	7.04	1.2%		0.6ft.	> 6.5ft.	Somewhat poorly drained	IIIw				64
MtA	Mitiwanga silt loam, 0 to 2 percent slopes	6.79	1.2%		1ft.	2.9ft. (Lithic bedrock)	Somewhat poorly drained	IIIw	110	33	50	65
MgB	Mahoning silt loam, 2 to 6 percent slopes	1.25	0.2%		0.6ft.	> 6.5ft.	Somewhat poorly drained	IIIe				62
Ln	Lorain silty clay loam	1.17	0.2%		0ft.	> 6.5ft.	Very poorly drained	IIIw	123	37	54	73
Weighted Average									4.2	1.3	1.9	63.6

*eftog PI: Obtained from the NRCS eFOTG (<http://efotg.sc.egov.usda.gov>)

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.