



# Soils Map



State: **Indiana**  
 County: **Pulaski**  
 Location: **5-29N-1W**  
 Township: **Van Buren**  
 Acres: **187.13**  
 Date: **2/7/2020**



Soils data provided by USDA and NRCS.

| Archived Soils Ending 1/21/2012 |   |       |                  |                      |             |   |                         |                  |       |          |              |
|---------------------------------|---|-------|------------------|----------------------|-------------|---|-------------------------|------------------|-------|----------|--------------|
| Code                            | Soil Description  | Acres | Percent of field | Non-Irr Class Legend | Water Table | Restrictive Layer                                     | Soil Drainage           | Non-Irr Class *c | Cor n | Soybeans | Winter wheat |
| BuuA                            | Brookston loam, 0 to 1 percent slopes                   | 65.79 | 35.2%            |                      | 0ft.        | > 6.5ft.  | Poorly drained          | IIw              | 173   | 49       | 70           |
| GmnA                            | Goodell-Gilford fine sandy loams, 0 to 1 percent slopes | 30.08 | 16.1%            |                      | 0.2ft.      | > 6.5ft.  | Poorly drained          | IIw              | 150   | 33       | 60           |
| BswA                            | Brems-Morocco loamy fine sands, 0 to 1 percent slopes   | 16.28 | 8.7%             |                      | 2.5ft.      | > 6.5ft.  | Moderately well drained | IVs              | 98    | 31       | 44           |
| CuyA                            | Crosier fine sandy loam, 0 to 1 percent slopes          | 13.14 | 7.0%             |                      | 1.2ft.      | 3.2ft. (Densic material)                              | Somewhat poorly drained | IIw              | 151   | 49       | 68           |
| MtpA                            | Moon-Selfridge complex, 0 to 1 percent slopes           | 12.66 | 6.8%             |                      | 2.2ft.      | > 6.5ft.  | Moderately well drained | IIIIs            | 131   | 44       | 60           |
| ScuA                            | Sebewa loam, 0 to 1 percent slopes                      | 11.67 | 6.2%             |                      | 0ft.        | 3ft. (Strongly contrasting textural stratification)   | Poorly drained          | IIw              | 133   | 35       | 54           |
| Mgza                            | Maumee-Gumz complex, 0 to 1 percent slopes              | 8.32  | 4.4%             |                      | 0.2ft.      | > 6.5ft.  | Poorly drained          | IIlw             | 139   | 30       | 56           |
| OacB                            | Oakville-Denham fine sands, 1 to 5 percent slopes       | 7.80  | 4.2%             |                      | 3ft.        | > 6.5ft.  | Excessively drained     | IVs              | 84    | 30       | 38           |
| MupA                            | Morocco loamy fine sand, 0 to 1 percent slopes          | 5.74  | 3.1%             |                      | 0.5ft.      | > 6.5ft.  | Somewhat poorly drained | IVs              | 102   | 29       | 46           |
| BwfA                            | Budd-Brady fine sandy loams, 0 to 1 percent slopes      | 4.02  | 2.1%             |                      | 0.5ft.      | 4.6ft. (Strongly contrasting textural stratification) | Somewhat poorly drained | IIw              | 126   | 36       | 57           |
| HtbAU                           | Houghton muck, undrained, 0 to 1 percent slopes         | 3.36  | 1.8%             |                      | 0ft.        | > 6.5ft.  | Very poorly drained     | Vw               |       |          |              |
| Mlwb                            | Metee-Moon loamy sands, 1 to 5 percent slopes           | 2.58  | 1.4%             |                      | 2.2ft.      | > 6.5ft.  | Well drained            | IIIe             | 122   | 43       | 58           |
| WpaA                            | Winamac-Bronson fine sandy loams, 0 to 1 percent slopes | 1.90  | 1.0%             |                      | 1.5ft.      | > 6.5ft.  | Moderately well drained | IIIs             | 127   | 43       | 57           |



|                         |   |      |      |  |          |          |                         |      |            |             |             |
|-------------------------|---|------|------|--|----------|----------|-------------------------|------|------------|-------------|-------------|
| ApuAN                   | Antung muck, drained, 0 to 1 percent slopes             | 1.89 | 1.0% |  | 0.2ft.   | > 6.5ft. | Very poorly drained     | Illw | 145        | 37          | 59          |
| CjfC                    | Chelsea fine sand, 5 to 12 percent slopes               | 1.35 | 0.7% |  | > 6.5ft. | > 6.5ft. | Excessively drained     | Vle  | 76         | 27          | 35          |
| SdzcB                   | Selfridge-Brems loamy fine sands, 1 to 4 percent slopes | 0.46 | 0.2% |  | 2.5ft.   | > 6.5ft. | Somewhat poorly drained | Illw | 116        | 37          | 53          |
| OaeC                    | Oakville fine sand, 5 to 12 percent slopes              | 0.09 | 0.0% |  | > 6.5ft. | > 6.5ft. | Excessively drained     | Vls  | 73         | 26          | 33          |
| <b>Weighted Average</b> |   |      |      |  |          |          |                         |      | <b>142</b> | <b>39.8</b> | <b>59.4</b> |

\*c: Using Capabilities Class Dominant Condition Aggregation Method  
Soils data provided by USDA and NRCS.